





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07B 53/00, 57/00, C07F 7/18, C07D 307/28, 309/18, C07F 11/00		A1	(11) International Publication Number: WO 00/02834
			(43) International Publication Date: 20 January 2000 (20.01.00)
(21) International Application Number: PCT/US99/15816			(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
(22) International Filing Date: 13 July 1999 (13.07.99)			
(30) Priority Data: 09/114,692 13 July 1998 (13.07.98) US			
(71) Applicant (for all designated States except US): MASSACHUSETTS INSTITUTE OF TECHNOLOGY [US/US]; 77 Massachusetts Avenue, Cambridge, MA 02139 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): SCHROCK, Richard, R. [US/US]; 15 Cabot Street, Winchester, MA 01890 (US). HOVEYDA, Amir, H. [US/US]; 20 Preble Gardens Road, Belmont, MA 02178 (US).			
(74) Agent: OYER, Timothy, J.; Wolf, Greenfield & Sacks, P.C., 600 Atlantic Avenue, Boston, MA 02210 (US).			Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

(54) Title: ASYMMETRIC METATHESIS REACTIONS INVOLVING ACHIRAL AND MESO SUBSTRATES

(57) Abstract

A composition and method for the catalytic conversion of a racemic mixture of dienes to a cyclic olefin by a ring-closing metathesis (RCM) reaction are disclosed. The composition, a transition metal complex with an M=C reaction site, contains a bidentate dialkoxide of at least 80 % optical purity. Because the M=C reaction site is of a sufficient shape specificity, conferred in part by the dialkoxide of sufficient rigidity and a M=N-R¹ site, reacting the composition with a mixture of two enantiomeric dienes results in an olefin metathesis product that has at least a 50 % enantiomeric excess of one enantiomer in the mixture. A method is also provided for reacting a composition with a racemic diene mixture to generate a metathesis product that has an enantiomeric excess of at least 50 %. Methods are also provided for catalytic enantioselective desymmetrization. One method involves an olefin metathesis reaction with a molecular substrate having a plane of symmetry to form a product free of a plane of symmetry. Another method provides a desymmetrization reaction to occur in the absence of solvent. A method for producing quaternary carbon centers through a desymmetrization reaction is also described.

**1a:** R³ = *t*-Bu; R¹ = R² = *i*-Pr**1b:** R³ = *t*-Bu; R¹ = R² = Me**1c:** R³ = adamantyl; R¹ = R² = *i*-Pr**1d:** R³ = adamantyl; R¹ = R² = Me**1e:** R³ = ethyl; R¹ = R² = *i*-Pr**1f:** R³ = ethyl; R¹ = R² = Me**2a:** R³ = 2,4,6-tri(*i*-propyl)phenyl;
R¹ = R² = *i*-Pr;**2b:** R³ = 2,4,6-tri(*i*-propyl)phenyl;
R¹ = R² = Me**2c:** R³ = *t*-Bu; R¹ = R² = *i*-Pr;
R⁴ = *t*-Bu**2d:** R³ = *t*-Bu; R¹ = R² = Me;
R⁴ = *t*-Bu**2e:** R³ = Ph; R¹ = R² = *i*-Pr; R⁴ = H**2f:** R³ = Ph; R¹ = R² = Me; R⁴ = H